[78]

XVIII. A Catalogue of the Fifty Plants from Chelsea Garden, presented to the Royal Society, by the worshipful Company of Apothecaries, for the Year 1754, pursuant to the Direction of Sir Hans Sloane, Baronet, Med. Reg. & Soc. Reg. nuper Præses, by John Wilmer, M. D. Societat. Pharmaceut. Lond. Socius, Hort. Chels. Præsectus & Præsector Botan.

Read April 10, 1601 A Ntirrhinum Siculum Linariæ folio, niveo flore, Tilli.

1602 Calcalia caule herbaceo foliis hastato-sagittatis denticulatis, petiolis superne dilatatis. Hort. Upsal. 254.

1603 Calamintha Hispanica frutescens Mari folio.

Inst. R. H. 194.

Melissa fruticosa ramis attenuatis virgatis soliis subtus tomentosis. Linn. Sp. Plant. 593.

1604 Calendula foliis obverse ovatis denticulatis, caule fruticoso perenni. Fl. Leyd. Prod. 531.

1605 Campanula hortensis rapunculi radice repente.M. H. 2. 460.

Campanula repens flore minore cæruleo. J. B. 2. 706.

1606 Caryophyllus minimus muralis. C. B. 211. Lychnis minima muralis. M. H. 2. 547.

1607 Cephalanthus foliis oppositis ternisque. Fl. Virg. 15.

Scabiosa

4

[79]

Scabiosa dendroides Americana ternis foliis caulem ambientibus, floribus ochro-leucis. Pluk. Alm. 336.

1608 Cerinthe quorundam major, fl. ex rubro-pur-

purascente. J. B. 3. 603.

1609 Chenopodium Lini folio villoso. T. 506. Linaria scoparia. C. B. P. 212.

1610 Clematis Hispanica surrecta altera et humilior, fl. albicante. H. R. Par.

1611 Clymenum Hispanicum flore vario, filiqua articulata. T. 396.

Lathyrus vicioides vexillo rubro, petalis roftrum ambientibus cæruleis. M. H. 2.50.

1612 Cochlearia altissima Glasti folio. T. 215. Lepidium. C. B. 96.

1613 Conyza latifolia viscosa suaveolens slore aureo, ex Gallo-provincia. Tourn. 455.

Conyza Pyrenaica foliis Primulæ veris. Par.

Bat. 127.

1614 Coreopsis foliis linearibus integerrimis caule erecto. Fl. Virg. 18.

1615 Corispermum Hyssopisolium. D. Justieu, Ac.

Reg. Sc. 1712.

1616 Digitalis Hispanica purpurea minor. T. 165. Digitalis Verbasci solio purpurea minor perennis Hispanica. Bar. Icon. 1183.

1617 Dipsacus folio laciniato. C. B. P. 385.

2618 Dodartia foliis lanceolatis ferratis, floribus spicatis. Linaria Bellidis folio. C. B. P. 212. Prod. 106.

1619 Fabago triphylla et tetraphylla flore tetrapetalo, fructu membranaceo quadrangulari. Burm. pl. Afric. 7. Tab. 3.

Zigo-

Zigophyllum capfularum angulis compresso membranaceis. Linn. Sp. Plant. 383.

1620 Geranium Anemones folio. Par. Bat. 178.

1621 Heliotropium Canariense arborescens, fol. Scorodoniæ. H. Am. 129.

1622 Hermannia fol. Ibisci hirsuto, slore primo luteo, deinde purpuro-rubello. An Hermannia frutescens folio oblongo molli cordato hirsuto. Boerh. Ind. Alt.

1623 Hesperis lutea filiquis strictissimis. Tourn. 222. Draba lutea quibusdam. J. B. 2. 870.

1624 Hieracium Echioides capitulis Cardui benedicti majus Arabicum folio finuato. Hort. Edinburg.

1625 Hieracium Pyrenaicum longifolium amplexi-

caule. T. 472.

1626 Jasminoides aculeatum humile Halimi minoris folio, slore majore violaceo, fructu crassiore per maturitatem slavescente. Michel. Nov. Gen. 224.

1627 Ketmia Ægyptiaca Vitis folio, parvo flore.

Tourn.

1628 Lathyrus Tingitanus filiquis Orobi, fl. amplo

ruberrimo. M. H. 2. 55.

1629 Lotus leguminibus conjugatis membranaceo quadrangulis bracteis oblongo-ovatis. Linn. Sp. Plant. 774.

Lotus luteus filiqua angulosa. Boer. Ind. Alt.

2. p. 37.

1630 Lychnis segetum rubra foliis Persoliatæ. C. B.

204. Vaccaria. Dod. Pempt. 104.

1631 Marrubium Hispanicum calyce stellato et aculeato. I. R. H. 182. Alysson Galeni. Clus. Hist. 35.

[81]

1632 Medica major erectior, floribus purpurascentibus. Tourn. 410.

1622 Mespilus foliis ovatis serratis, pomis ovatis ex luteo virentibus nobis.

1624 Mespilus foliis ovatis, obtusis, crenatis, spinis longis tenuibus, fructu fingulari rotundo, umbilico foliaceo. An Mespilus Virginiana. Uvæ crispæ foliis. Pluk. Mant.

1635 Ornithogalum humile foliis ovatis sessilibus, floribus corymbosis, nobis.

1626 Osteospermum foliis oblongis acute dentatis,

caule fruticoso diffuso, nobis.

1627 Othonna foliis pinnatifidis laciniis linearibus parallelis. H. Cliff. 419. Jacobæa Africana frutescens foliis Absinthii umbelliseri incanis. H. Amst. 2. p. 137.

1638 Pelecinus vulgaris. T. 417. Astragalus purpureus annuus peregrinus, filiquis utriusque

ferræ fimilibus. M. Hist. 2. 107.

1639 Phyllis stipulis dentatis. Flor. Leyd. Prod. 92. Valerianella Canariens. frutescens simpla nobla dicta. Hort. Eltham 405.

1640 Plantago angustifolia paniculis Lagopi. C. B.

374.

1641 Ruyschiana floribus verticillatis, nobis.

1642 Samolus Valerandi. J. B. 3. 791. Anagallis aquatica folio rotundo non crenato. C. B. P. 252.

1642 Scabiosa stellata folio laciniato major. Tourn.

1644 Solanum spiniferum frutescens spinis igneis Americanum. Pluk. Phyt. 225, fig. 5.

1645 Telephium Dioscoridis Imperati.

Telephium repens folio non deciduo. C.B.P. 287.

1646 Thalictrum minus alterum Parifiensium foliis crassioribus et lucidis. H. R. Par.

Tithymalus maritimus. C. B. P. 291. Tithimalus paralius. J. B. 3. 674.

1648 Trifolium Bitumen redolens. C. B. 327.

1649 Trifolium Bitumen redolens angustifolium. Boer. Ind. Alt. 2. p. 32.

> Trifolium bituminosum arboreum angustifolium ac sempervirens. Hort. Cath.

1650 Virga aurea Canadensis foliis carnosis non serratis latioribus. Hist. Oxon.

XIX. A Letter to the Right Honourable George Earl of Macclesfield, President of the Royal Society, on the Advantage of taking the Mean of a Number of Observations, in practical Astronomy: By T. Simpson, F. R. S.

My Lord,

Read April 10, T is well known to your Lordship, 1755. That the method practifed by astronomers, in order to diminish the errors arising from the impersections of instruments, and of the organs of sense, by taking the Mean of several observations, has not been so generally received, but that some persons, of considerable note, have been of opinion, and even publickly maintained, that one single observation,